

QCD description of high order factorial moments in quark and gluon jets ^a

SERGIO LUPIA

*Max-Planck-Institut für Physik
(Werner-Heisenberg-Institut)
Föhringer Ring 6, D-80805 München, Germany*

Abstract


The complete QCD evolution equation for factorial moments in quark and gluon jets is numerically solved with initial conditions at threshold by fully taking into account the energy-momentum conservation law. Within the picture of Local Parton Hadron Duality, the perturbative QCD predictions can successfully describe the available experimental data.

^ato be published in the Proceedings of the XXXIIIrd Rencontres de Moriond, “QCD and high energy interactions”, Les Arcs, France, March 21-28, 1998

QCD DESCRIPTION OF HIGH ORDER FACTORIAL MOMENTS IN QUARK AND GLUON JETS

SERGIO LUPIA

*Max-Planck-Institut für Physik (Werner-Heisenberg-Institut),
Föhringer Ring 6, 80805 München, Germany*



The complete QCD evolution equation for factorial moments in quark and gluon jets is numerically solved with initial conditions at threshold by fully taking into account the energy-momentum conservation law. Within the picture of Local Parton Hadron Duality, the perturbative QCD predictions can successfully describe the available experimental data.

1 Introduction

An interesting approach for describing multiparticle production in high energy jets is provided by the analytical perturbative approach¹. In this framework, the perturbative evolution is described in terms of a parton cascade which is evolved down to small scales of a few hundred MeV for the transverse momentum, and then partonic predictions are compared to hadronic observables, according to the notion of local parton hadron duality (LPHD)². In the description of the partonic cascade QCD coherence is included as well as the running of the coupling α_s at one-loop level and energy-momentum conservation effects. This approach has been shown to be phenomenologically rather successful for a large set of observables³, and in particular for jet and particle multiplicity⁴. However, recent experimental analyses⁵ have found a quantitative discrepancy between data on factorial cumulants and factorial moments of the multiplicity distribution in single quark and gluon jets and theoretical predictions, as given in⁶, where only the leading part of the energy-momentum conservation effects have been taken into account.

Here we report the result of a new analysis⁷, where the complete MLLA evolution equation of QCD for factorial moments of any order with the full inclusion of energy-momentum conservation effects have been numerically solved. After LPHD, the analytical perturbative approach is then shown to be able to describe quantitatively the available experimental data on high order factorial moments and cumulants in single quark and gluon jets.

q	$F_q^{(q)}$ Quark jet		$F_g^{(q)}$ Gluon jet	
	Exp ⁵	Theory	Exp ⁵	Theory
2	$1.0820 \pm 0.0006 \pm 0.0046$	1.080	$1.023 \pm 0.008 \pm 0.011$	1.026
3	$1.275 \pm 0.002 \pm 0.017$	1.265	$1.071 \pm 0.026 \pm 0.034$	1.078
4	$1.627 \pm 0.005 \pm 0.042$	1.600	$1.146 \pm 0.059 \pm 0.074$	1.157
5	$2.274 \pm 0.014 \pm 0.093$	2.168	$1.25 \pm 0.11 \pm 0.13$	1.268

Table 1: Experimental data for the normalized factorial moments of order $q = 2, \dots, 5$ in single quark and gluon jets of 45.6 and 41.8 GeV respectively are compared with our theoretical predictions.

2 Results

We have considered the complete MLLA evolution equations for the generating function¹ and derived the corresponding system of coupled evolution equations for unnormalized factorial moments $\tilde{F}^{(q)}$. We have then numerically solved this system by imposing boundary conditions at threshold, i.e., $\mathcal{N}_g = \mathcal{N}_q = 1$ and $\tilde{F}_g^{(q)} = \tilde{F}_q^{(q)} = 0$ for $q > 1$.

The three free parameters, i.e., the infrared cutoff Q_0 , the QCD scale Λ and the normalization parameter K_{all} , have been kept equal to the values fixed in⁴ to describe the jet and particle multiplicities:

$$K_{all} = 1, \quad \Lambda = 500 \text{ MeV} \quad Q_0 = 0.507 \text{ MeV} . \quad (1)$$

A very good quantitative agreement with the experimental data of⁵ is achieved, contrary to previous comparisons⁵ with approximate predictions⁶.

We refer to⁷ for more details on this new analysis.

3 Conclusions

We have shown that the analytical perturbative approach, based on perturbative QCD description of the partonic evolution plus Local Parton Hadron Duality as hadronization prescription, can quantitatively describe experimental data on multiparticle correlations in single quark and gluon jets, if the energy-momentum conservation law is correctly taken into account.

Acknowledgments

I would like to thank Wolfgang Ochs for many fruitful discussions and suggestions.

References

1. Yu. L. Dokshitzer, V. A. Khoze, A. H. Mueller and S. I. Troyan, Rev. Mod. Phys. **60** (1988) 373; ‘Basics of Perturbative QCD”, ed. J. Tran Thanh Van, Editions Frontières, Gif-sur-Yvette, 1991.
2. Ya. I. Azimov, Yu. L. Dokshitzer, V. A. Khoze and S. I. Troyan, Z. Phys. **C27** (1985) 65 and **C31** (1986) 213.
3. V.A. Khoze and W. Ochs, Int. J. Mod. Phys. **A12** (1997) 2949.
4. S. Lupia and W. Ochs, Phys. Lett. **B418** (1998) 214.
5. OPAL Coll., K. Akerstaff et al., Eur. Phys. Journ. **C1** (1998) 479.
6. I. M. Dremin and V. A. Nechitailo, Mod. Phys. Lett. **A9** (1994) 1471.
7. S. Lupia, “Perturbative QCD description of multiparticle correlations in quark and gluon jets”, in preparation